

On the confidence interval for the parameter of Poisson distribution

S.I. Bityukov^{a,1} N.V. Krasnikov^b, V.A. Taperechkina^c

^aDivision of Experimental Physics, Institute for High Energy Physics, Protvino, Moscow Region, Russia

^bDivision of Quantum Field Theory, Institute for Nuclear Research RAS, Moscow, Russia

^cMathematics and Computer Science Department, Moscow State Academy of Instrument Engineering and Computer Science, Serpukhov, Moscow Region, Russia

Abstract

The possibility of construction of continuous analogue of Poisson distribution with the search of bounds of confidence intervals for parameter of Poisson distribution is discussed. Also, in the article is shown that the true value of a parameter of Poisson distribution for the observed value $\hat{x}$ has Gamma distribution with the scale parameter, which is equal to one, and the shape parameter, which is equal to $\hat{x}$. The results of numerical construction of confidence intervals are presented.

Keywords: statistics, confidence intervals, Poisson distribution, Gamma distribution, sample.

1 Introduction

In article [1] the unified approach to the construction of confidence intervals and confidence limits for a signal with a background presence, in particular for Poisson distributions, is proposed. The method is widely used for the presentation of physical results [2] though a number of investigators criticize this approach [3] (in particular, this approach avoids a violation of the coverage principle). Series of Workshops on Confidence Limits has been held in CERN and Fermilab. At these meetings demands for properties of constructed confidence intervals and confidence limits have been formulated [4]. On the other hand, the results of experiments often give noninteger values of a number of observed events (for example, after background subtraction [5])

¹Corresponding author

Email addresses: bityukov@mx.ihep.su, Serguei.Bitoukov@cern.ch

when a Poisson distribution take place. That is why there is a necessity to search a continuous analogue of Poisson distribution. The present work offers some generalization of Poisson distribution for continuous case. The generalization given here allows to construct confidence intervals and confidence limits for Poisson distribution parameter both for integer and real values of a number of observed events, using conventional methods. More than, the supposition about continuous of some function $f(x, \lambda)$ described below allows to use a Gamma distribution for construction of confidence intervals and confidence limits of Poisson distribution parameter. In present article we consider only the construction of confidence intervals.

In the Section 2 the generalization of Poisson distribution for the continuous case is introduced. An example of confidence intervals construction for the parameter of analogue of Poisson distribution is given in the Section 3. In the Section 4 the results of construction of confidence intervals having the minimal length for the parameter of Poisson distribution using a Gamma distribution are discussed. The main results of the article are formulated in the Conclusion.

2 The Generalization of Discrete Poisson Distribution for the Continuous Case

Let us have a random value ξ , taking values from the set of numbers $x \in X$. Let us consider two-dimensional function

$$f(x, \lambda) = \frac{\lambda^x}{x!} e^{-\lambda}, \quad (1)$$

where $x \geq 0$ $\lambda > 0$. Assume, that set X includes only integer numbers, then discrete function $f(x, \lambda)$ describes distribution of probabilities for Poisson distribution with the parameter λ and a random variable x

Let us rewrite the density of Gamma distribution using unconventional notation

$$f(x, a, \lambda) = \frac{a^{x+1}}{\Gamma(x+1)} e^{-a\lambda} \lambda^x, \quad (2)$$

where a is a scale parameter, $x > -1$ is a shape parameter² and $\lambda > 0$ is a random variable. Here the quantities of x and λ take values from the set of

²A conventional notation for the shape parameter is [6] $r = x + 1$, $r > 0$.

real numbers. Let us set $a = 1$ and denote $x! = \Gamma(x + 1)$, then a continuous function

$$f(x, \lambda) = \frac{\lambda^x}{x!} e^{-\lambda}, \quad \lambda > 0, \quad x > -1 \quad (3)$$

is the density of Gamma distribution with the scale parameter $a = 1$.

To anticipate a little, it is indicative of the Gamma distribution of parameter λ for the Poisson distribution in case of observed value $x = \hat{x}$.

Figure 1 shows the surface described by the function $f(x, \lambda)$. Smooth behaviour of this function along x and λ (see Fig.2) allows to assume that there is such a function $l(\lambda) > -1$, that

$$\int_{l(\lambda)}^{\infty} f(x, \lambda) dx = 1 \quad (4)$$

for given value of λ . It means that in this way we introduce continued analogue of Poisson distribution with the probability density $f(x, \lambda) = \frac{\lambda^x}{x!} e^{-\lambda}$ over the area of the function definition, i.e. for $x \geq l(\lambda)$ and $\lambda > 0$. The values of the function $f(x, \lambda)$ for integer x coincide with corresponding magnitudes in the probabilities distribution of discrete Poisson distribution. Dependences of the values of function $l(\lambda)$, the means and the variances for the suggested distribution on λ have been calculated by using programme DGQUAD from the library CERNLIB [7] and the results are presented in Table 1. This Table shows that series of properties of Poisson distribution ($E\xi = \lambda, D\xi = \lambda$) take place only if the value of the parameter $\lambda > 3$.

It is appropriate at this point to say that

$$\int_0^{\infty} f(x, \lambda) dx = \int_0^{\infty} \frac{\lambda^x e^{-\lambda}}{\Gamma(x + 1)} dx = e^{-\lambda} \nu(\lambda). \quad (5)$$

The function

$$\nu(\lambda) = \int_0^{\infty} \frac{\lambda^x}{\Gamma(x + 1)} dx \quad (6)$$

is well known and, according to ref. [8],

$$\nu(\lambda) = \sum_{n=-N}^{\infty} \frac{\lambda^n}{\Gamma(n + 1)} + O(|\lambda|^{-N-0.5}) = e^{\lambda} + O(|\lambda|^{-N}) \quad (7)$$

if $\lambda \rightarrow \infty$, $|\arg \lambda| \leq \frac{\pi}{2}$ for any integer N . Nevertheless we have to use the function $l(\lambda)$ in our calculations in Section 3. We consider it as a mathematical trick to illustrate a possibility of construction of confidence intervals by numerically in the case of real value $\hat{x}$.

In principle, we can numerically transforms the function $f(x, \lambda)$ in the interval $x \in (0, 1)$ so that

$$\int_0^\infty f(x, \lambda) dx = 1, \quad E\xi = \int_0^\infty x f(x, \lambda) dx = \lambda,$$

$$D\xi = \int_0^\infty (x - E\xi)^2 f(x, \lambda) dx = \lambda$$

for any λ . In this case we can construct confidence intervals without introducing of $l(\lambda)$.

In Section 3 assumption about continuous of the function $f(x, \lambda)$ along the variable x are used for construction of confidence intervals of a parameter λ for any observed $\hat{x}$.

Let us construct a central confidence intervals for the continued analogue of Poisson distribution using function $l(\lambda)$.

3 The Construction of the Confidence Intervals for Continued Analogue of Poisson Distribution.

Assume that in the experiment with the fixed integral luminosity (i.e. a process under study may be considered as a homogeneous process during given time) the $\hat{x}$ events of some Poisson process were observed. It means that we have an experimental estimation $\hat{\lambda}(\hat{x})$ of the parameter λ of Poisson distribution. We have to construct a confidence interval $(\hat{\lambda}_1(\hat{x}), \hat{\lambda}_2(\hat{x}))$, covering the true value of the parameter λ of the distribution under study with confidence level $1 - \alpha$, where α is a significance level. It is known from the theory of statistics [9], that the mean value of a sample of data is an unbiased estimation of mean of distribution under study. In our case the sample consists of one observation $\hat{x}$. For the discrete Poisson distribution the mean coincides with the estimation of parameter value, i.e. $\hat{\lambda} = \hat{x}$. This is not true for a small value of λ in the considered case (see Table 1). That is why in order to find the estimation of $\hat{\lambda}(\hat{x})$ for small value $\hat{x}$ there is necessary to introduce correction in accordance with Table 1. Let us construct the central confidence intervals using conventional method assuming that

$$\int_{\hat{x}}^\infty f(x, \hat{\lambda}_1) dx = \frac{\alpha}{2} \quad (8)$$

for the lower bound $\hat{\lambda}_1$ and

$$\int_{l(\hat{\lambda}_2)}^{\hat{x}} f(x, \hat{\lambda}_2) dx = \frac{\alpha}{2} \quad (9)$$

for the upper bound $\hat{\lambda}_2$ of confidence interval.

Figure 3 shows the introduced in the Section 2 distributions with parameters defined by the bounds of confidence interval ($\hat{\lambda}_1 = 1.638, \hat{\lambda}_2 = 8.493$) for the case $\hat{x} = \hat{\lambda} = 4$ and the Gamma distribution with parameters $a = 1, x = \hat{x} = 4$. The association between the confidence interval and the Gamma distribution is seen from this Figure. The bounds of confidence interval with 90% confidence level for the parameter of continued analogue of Poisson distribution for different observed values $\hat{x}$ (first column) were calculated and are given in the second column of the Table 2. It is necessary to notice that the confidence level of the constructed confidence intervals always coincides exactly with the required confidence level. As it results from Table 2 that the suggested approach allows to construct confidence intervals for any real and integer values of the observed number of events in the case of the values of parameter $\lambda > 3$. The Table 2 shows that the left bound of central confidence intervals is not equal to zero for small $\hat{x}$, what is not suitable.

Also note that 90% of the area of Gamma distributions with parameter $x = \hat{x}$ are contained inside the constructed 90% confidence intervals for observed value $\hat{x}$. However, for small values of $\hat{x}$ we have got values of the area close to 88%, i.e. less than 90%.

4 Shortest Confidence Intervals for Parameter of Poisson Distribution.

As is follow from the formula (3) for $f(x, \lambda)$ (see Fig.3) the true value of λ of Poisson distribution for the observed value $\hat{x}$ has Gamma distribution (at least both for integer $x \geq 0$ if we consider pure Poisson distribution and for real $x \geq 0$ if we consider analogue of Poisson distribution) with the parameters $a = 1$ and $x = \hat{x}$, i.e.

$$P(\lambda|\hat{x}) = \frac{\lambda^{\hat{x}}}{\hat{x}!} e^{-\lambda}. \quad (10)$$

This supposition allows to choose confidence interval of a minimum length from all possible confidence intervals of given confidence level without violation of the coverage principle. The bounds of minimum length area, con-

taining 90% of the corresponding Gamma distribution square, were found numerically both for integer value of $\hat{x}$ and for real value of $\hat{x}$. We took into account that $x = \hat{x}$ and found the shortest 90% confidence interval for the parameter of Poisson distribution. The results are presented in third column of the Table 2. For comparison with the results of conventional procedure [2] of finding confidence intervals, the results of calculations of confidence intervals for integer value of $\hat{x}$ [1] are adduced in the Table 2. By this means confidence intervals, got using Gamma distribution, may be used for real values of $\hat{x}$.

5 Conclusion

In the article the attempt of introducing of continued analogue of Poisson distribution for the construction of classical confidence intervals for the parameter λ of Poisson distribution is described. Also, in the article is shown that the true value of a parameter of Poisson distribution for the observed value $\hat{x}$ has Gamma distribution with the scale parameter $a = 1$ and the shape parameter $x = \hat{x}$. Two approaches (with using of function $l(\lambda)$ and with using of Gamma distribution) are considered. Confidence intervals for different integer and real values of number of observed events for Poisson process in the experiment with given integral luminosity are constructed. The second approach allows to construct any confidence intervals for a parameter of Poisson distribution.

Acknowledgments

We are grateful to V.A. Matveev, V.F. Obraztsov and Fred James for the interest to this work and for valuable comments. We are thankful to S.S. Bitjukov and V.A. Litvine for useful discussions. We would like to thank E.A. Medvedeva for the help in preparing the article. This work has been supported by RFFI grant 99-02-16956 and grant INTAS-CERN 377.

References

- [1] Feldman, G.J. and R.D. Cousins, Unified approach to the classical statistical analysis of small signal, *Phys.Rev.* **D57** 3873-3889 (1998).
- [2] Groom, D.E. et al., Review of particle physics, by J. Bartels, D. Haidt and A. Zichichi (Eds.), *Eur.Phys.J.* **C 3**, (1998) 198-199.
- [3] as an example, Zech, G., Classical and Bayesian Confidence Limits, in: F. James, L. Lyons, and Y. Perrin (Eds.), *Proc. of 1st Workshop on Confidence Limits*, CERN 2000-005, Geneva, Switzerland, (2000) 141-154.
- [4] James, F., Introduction and Statement of the Problem, in: F. James, L. Lyons, and Y. Perrin (Eds.), *Proc. of 1st Workshop on Confidence Limits*, CERN 2000-005, Geneva, Switzerland, (2000) 1-2.
- [5] Cronin-Hennessy, D. et al., Observation of $B \rightarrow K^\pm \pi^0$ and $B \rightarrow K^0 \pi^0$ and Evidence for $B \rightarrow \pi^+ \pi^-$. *Phys.Rev.Lett.D* **85** (2000) 515.
- [6] Kendall, Maurice G; Stuart, Alan; J. Keith Ord, *The advanced theory of statistics* . Vol. 1 *Distribution theory*. 6th ed. (Arnold, A member of the Hodder Headline Group, 338 Euston Road, London, 1994) 182.
- [7] CERNLIB, CERN PROGRAM LIBRARY, Short Writeups, (CERN, Geneva, Switzerland, Edition - June 1996)
- [8] Bateman H., *Higher Transcendental Functions*. Vol. 3, A. Erdélyi (Ed.); (McGraw-Hill Book Company Inc., 1955) 217-224.
- [9] as an example, *Handbook of Probability Theory and Mathematical Statistics (in Russian)*, V.S. Korolyuk (Ed.), (Kiev, "Naukova Dumka", 1978)

Table 1: The function $l(\lambda)$, mean and variance versus λ .

λ	$l(\lambda)$	mean ($E\xi$)	variance ($D\xi$)
0.001	-0.297	-0.138	0.024
0.002	-0.314	-0.137	0.029
0.005	-0.340	-0.130	0.040
0.010	-0.363	-0.120	0.052
0.020	-0.388	-0.100	0.071
0.050	-0.427	-0.051	0.113
0.100	-0.461	0.018	0.170
0.200	-0.498	0.142	0.272
0.300	-0.522	0.256	0.369
0.400	-0.539	0.365	0.464
0.500	-0.553	0.472	0.559
0.600	-0.564	0.577	0.653
0.700	-0.574	0.681	0.748
0.800	-0.582	0.785	0.844
0.900	-0.590	0.887	0.939
1.00	-0.597	0.989	1.035
1.50	-0.622	1.495	1.521
2.00	-0.639	1.998	2.012
2.50	-0.650	2.499	2.506
3.00	-0.656	3.000	3.003
3.50	-0.656	3.500	3.501
4.00	-0.647	4.000	3.999
4.50	-0.628	4.500	4.498
5.00	-0.593	5.000	4.997
5.50	-0.539	5.500	5.497
6.00	-0.466	6.000	5.996
6.50	-0.373	6.500	6.495
7.00	-0.262	7.000	6.995
7.50	-0.135	7.500	7.494
8.00	0.000	8.000	7.993
8.50	0.000	8.500	8.496
9.00	0.000	9.000	8.997
9.50	0.000	9.500	9.498
10.0	0.000	10.00	9.999

Table 2: 90% C.L. intervals for the Poisson signal mean λ for total events observed $\hat{x}$.

$\hat{x}$	bounds (Section 3)		bounds (Section 4)		bounds (ref[1])	
	$\hat{\lambda}_1$	$\hat{\lambda}_2$	$\hat{\lambda}_1$	$\hat{\lambda}_2$	$\hat{\lambda}_1$	$\hat{\lambda}_2$
0.000	0.121E-08	2.052	0.0	2.303	0.00	2.44
0.001	0.205E-08	2.054	0.0	2.304		
0.002	0.292E-08	2.056	0.0	2.306		
0.005	0.666E-08	2.061	0.0	2.311		
0.01	0.307E-07	2.076	0.0	2.320		
0.02	0.218E-06	2.098	0.0	2.337		
0.05	0.765E-05	2.166	1.66E-05	2.389		
0.10	0.137E-03	2.275	2.23E-05	2.474		
0.20	0.186E-02	2.490	6.65E-05	2.642		
0.30	0.696E-02	2.692	1.49E-04	2.806		
0.40	0.161E-01	2.891	2.60E-03	2.969		
0.50	0.295E-01	3.084	5.44E-03	3.129		
0.60	0.466E-01	3.269	1.35E-02	3.290		
0.70	0.673E-01	3.450	2.63E-02	3.452		
0.80	0.911E-01	3.629	4.04E-02	3.611		
0.90	0.1179	3.804	6.12E-02	3.773		
1.0	0.1473	3.977	8.49E-02	3.933	0.11	4.36
1.5	0.3257	4.800	0.2391	4.718		
2.0	0.5429	5.582	0.4410	5.479	0.53	5.91
2.5	0.7896	6.340	0.6760	6.220		
3.0	1.056	7.076	0.9284	6.937	1.10	7.42
3.5	1.340	7.792	1.219	7.660		
4.0	1.638	8.493	1.511	8.358	1.47	8.60
4.5	1.946	9.188	1.820	9.050		
5.0	2.264	9.869	2.120	9.714	1.84	9.99
5.5	2.590	10.55	2.453	10.39		
6.0	2.924	11.21	2.775	11.05	2.21	11.47
6.5	3.264	11.87	3.126	11.72		
7.0	3.609	12.53	3.473	12.38	3.56	12.53
7.5	3.961	13.18	3.808	13.01		
8.0	4.316	13.82	4.160	13.65	3.96	13.99
8.5	4.677	14.46	4.532	14.30		
9.0	5.041	15.10	4.905	14.95	4.36	15.30
9.5	5.406	15.73	5.252	15.56		
10.	5.779	16.36	5.640	16.21	5.50	16.50
20.	13.65	28.49	13.50	28.33	13.55	28.52




Figure 1: The behaviour of the function $f(x, \lambda)$ versus λ and x if $f(x, \lambda) < 1$.




Figure 2: Two-dimensional representation of the function $f(x, \lambda)$ versus λ and x for values $f(x, \lambda) < 1$.




Figure 3: The probability densities $f(x, \lambda)$ of continuous analogue Poisson distribution for λ 's determined by the confidence limits $\hat{\lambda}_1$ and $\hat{\lambda}_2$ in case of observed number of events $\hat{x} = 4$ and the probability density of Gamma distribution with parameters $a = 1$ and $x = \hat{x} = 4$.